

产品概述

产品名 (Product Name)	CD32-C rabbit Polyclonal Antibody
货号 (Catalog No.)	ATA26324
种类 (Category)	Primary antibodies
宿主 (Host)	Rabbit
反应种属 (Species specificity)	Human
应用实验 (Tested applications)	WB,ELISA
克隆性 (Clonality)	Polyclonal
偶连物 (Conjugation)	Unconjugated
免疫原 (Immunogen)	The antiserum was produced against synthesized peptide derived from human FCGR2C. AA range:251-300

产品性能

状态 (Form)	Liquid
存放条件 (Storage)	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 4 °C for frequent use. Store at -20 to -80 °C for twelve months from the date of receipt.
纯化方式 (Purity)	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

应用

Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

产品背景

caution:Has sometimes been attributed to correspond to FcR-IIb.,caution:Has sometimes been attributed to correspond to FcR-IIc.,disease:A chromosomal aberration involving FCGR2B is found in a follicular lymphoma. Translocation t(1;22)(q22;q11). The translocation leads to the hyperexpression of the receptor. This may play a role in the tumor progression.,domain:Contains 1 copy of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,domain:Contains an intracytoplasmic twice repeated motif referred as immunoreceptor tyrosine-based activator motif (ITAM). These motifs are involved in triggering cell activation upon receptors aggregation.,function: Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells.,function:

Receptor for the Fc region of complexed or aggregated immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells. Binding to this receptor results in down-modulation of previous state of cell activation triggered via antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Isoform IIB1 interacts with measles virus N protein. N protein is released in the blood following lysis of measles infected cells. This interaction presumably block inflammatory immune response. Interacts with INPP5D/SHIP1.,tissue specificity:Is the most broadly distributed Fc-gamma-receptor. Expressed in monocyte, neutrophils, macrophages, basophils, eosinophils, Langerhans cells, B-cells, platelets cells and placenta (endothelial cells). Not detected in natural killer cells.,tissue specificity:Isoform IIC1 is detected in monocytes, macrophages, polymorphonuclear cells and natural killer cells.,

