

产品概述

产品名 (Product Name)	CD33 rabbit Polyclonal Antibody
货号 (Catalog No.)	ATA26325
种类 (Category)	Primary antibodies
宿主 (Host)	Rabbit
反应种属 (Species specificity)	Human
应用实验 (Tested applications)	WB,ELISA
克隆性 (Clonality)	Polyclonal
偶连物 (Conjugation)	Unconjugated
免疫原 (Immunogen)	The antiserum was produced against synthesized peptide derived from the Internal region of human CD33. AA range:71-120

产品性能

状态 (Form)	Liquid
存放条件 (Storage)	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 4 °C for frequent use. Store at -20 to -80 °C for twelve months from the date of receipt.
纯化方式 (Purity)	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

应用

Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

产品背景

domain:Contains 2 copies of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,function:Putative adhesion molecule of myelomonocytic-derived cells that mediates sialic-acid dependent binding to cells. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic acids on the same cell surface. In the immune response, may act as an inhibitory receptor upon ligand induced tyrosine phosphorylation by recruiting cytoplasmic phosphatase(s) via their SH2 domain(s) that block signal transduction through dephosphorylation of signaling molecules. Induces apoptosis in acute myeloid leukemia (in vitro).,online information:Siglec-3,PTM:Phosphorylation of Tyr-340 is involved in binding to PTPN6 and PTPN11. Phosphorylation of Tyr-358 is involved in binding to PTPN6.,similarity:Belongs to the immunoglobulin superfamily. SIGLEC (sialic acid binding Ig-like lectin)

Catalog Number:ATA26325

CD33 rabbit Polyclonal Antibody

family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain.,subunit:Interacts with PTPN6/SHP-1 and PTPN11/SHP-2 upon phosphorylation.,tissue specificity:Monocytic /myeloid lineage cells.,

