

产品概述

产品名 (Product Name)	CD40 rabbit Polyclonal Antibody
货号 (Catalog No.)	ATA26355
种类 (Category)	Primary antibodies
宿主 (Host)	Rabbit
反应种属 (Species specificity)	Human,Mouse,Monkey
应用实验 (Tested applications)	WB,IHC-p,ELISA
克隆性 (Clonality)	Polyclonal
偶连物 (Conjugation)	Unconjugated
免疫原 (Immunogen)	The antiserum was produced against synthesized peptide derived from human CD40. AA range:228-277

产品性能

状态 (Form)	Liquid
存放条件 (Storage)	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 4 °C for frequent use. Store at -20 to -80 °C for twelve months from the date of receipt.
纯化方式 (Purity)	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

应用

Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

产品背景

CD40 molecule(CD40) Homo sapiens This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIG